Serial No. 10/659,598

Listing of Claims:

1-72 (canceled)

- 73. (Currently amended) A method of treating acne comprising topically administering to a mammal in need of treatment therefor an effective amount of a composition comprising (a) tretinoin and (b) a nondenatured soybean extract which has not been denatured having trypsin-inhibiting activity.
- 74. (Previously presented) A method according to claim 73 wherein said nondenatured soybean extract is selected from the group consisting of soybean milk and soybean paste.
- 75. (Previously presented) A method of treating acne comprising topically administering to a mammal in need of treatment therefor an effective amount of a composition comprising (a) retinol and (b) a nondenatured sovbean extract having trynsin-inhibiting activity.
- 76. (Previously presented) A method according to claim 75 wherein said nondenatured soybean extract is selected from the group consisting of soybean milk and soybean paste.
- 77. (Previously presented) A method according to claim 73 wherein said nondenatured soybean extract is soybean powder.
- 78. (Previously presented) A method according to claim 75 wherein said nondenatured soybean extract is soybean powder.
- 79. (Previously presented) A method of treating acne comprising topically administering to a mammal in need of treatment therefor an effective amount of a composition comprising (a) about 0.01 to about 0.3 weight percent of tretinoin and (b) about 0.01 to about 50 weight percent of a nondenatured soybean extract having trypsin-inhibiting activity.
- 80. (Previously presented) A method according to claim 79 wherein said nondenatured soybean extract is a soy powder.

Serial No. 10/659,598

- 81. (Previously presented) A method according to claim 80 wherein said soy powder is present in an amount of about 0.05 to about 20 weight percent.
- 82. (Previously presented) A method according to claim 81 wherein said soy powder is present in an amount of about 0.5 to about 5 weight percent.
- 83. (Previously presented) A method of treating acne comprising topically administering to a mammal in need of treatment therefor an effective amount of a composition comprising (a) about 0.1 to about 1 weight percent of retinol and (b) about 0.01 to about 50 weight percent of a nondenatured soybean extract having trypsin-inhibiting activity.
- 84. (Previously presented) A method according to claim 83 wherein said nondenatured soybean extract is a soy powder.
- 85. (Previously presented) A method according to claim 84 wherein said soy powder is present in an amount of about 0.05 to about 20 weight percent.
- 86. (Previously presented) A method according to claim 85 wherein said soy powder is present in an amount of about 0.5 to about 5 weight percent.